



**MALCOLM FRAZER**  
GYNAECOLOGIST  
UROGYNAECOLOGIST

**Assoc/Prof Malcolm Frazer**  
MB ChB MD FRCOG FRANZCOG CU  
Gynaecologist/Urogynaecologist

140 Ashmore Road, Benowa 4217  
Gold Coast and Pindara Hospitals Griffith and Bond University, Gold Coast, Australia

**Phone:** +61 (0) 755649300  
**Fax:** +61 (0) 755649400  
**Email:** [info@malcolmfrazer.com.au](mailto:info@malcolmfrazer.com.au)

---

## **DIVISION OF SUBURETHRAL SLING (SUS) OR TRIMMING OF MESH**

**You may have developed voiding dysfunction or have erosion of the tape and division of your sling has been suggested.**

**This is performed as a day surgery.**

### **DIVISION OF SUS:**

Division of the tape is performed under a general anaesthesia in the operating theatre. This allows better view of the precise area that is to be divided or removed. It is usually a very small area of the tape being exposed or you have developed problems emptying your bladder and being on the anterior wall of the vagina it is very difficult to perform this in an outpatient setting.

### **WHAT DOES IT INVOLVE?**

There is a very small incision on the front wall of the vagina and once the tape is trimmed or divided 1 or 2 small stitches are used to close the incision. These will dissolve in usually 2 to 3 weeks. You may experience some vaginal bleeding during this period of time but it should only be spotting.

### **TRIMMING OF VAGINAL MESH:**

Trimming of the vaginal mesh is performed under a general anaesthesia in the operating theatre. This allows better access to the area of mesh that needs to be trimmed.

### **WHAT DOES IT INVOLVE?**

A small incision is made in the area of mesh rejection and the mesh is trimmed. 1 to 2 stitches are used depending on the incision. These will dissolve in usually 2 to 3 weeks. You may experience some vaginal bleeding during this period of time but it should only be spotting.

### **RECOVERY:**

If this procedure is being performed within 6 weeks post operatively you need to continue with the normal post op recommendations.

If this procedure is being performed after 6 weeks you will be able to return to work in 1 to 2 days depending on your recovery from the anaesthetic. There are no restrictions on normal everyday activities but you should abstain from intercourse or the use of tampons for 2 weeks to allow the vaginal incision to heal and stitches to dissolve.

This procedure should not interfere with the result of your previous surgery and you will be reviewed in the clinic 4 to 6 weeks after this procedure.